

For Research Use Only

# Anti-Human PD-1/CD279 Rabbit Recombinant Antibody

Catalog Number: 98068-1-RR



## Basic Information

Catalog Number:

98068-1-RR

Concentration:

100ug, 1000 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC074740

GeneID (NCBI):

5133

UNIPROT ID:

Q15116

Full Name:

programmed cell death 1

Calculated MW:

288 aa, 32 kDa

Purification Method:

Protein A purification

CloneNo.:

240724G11

Recommended Dilutions:

IHC 1:500-1:2000

## Applications

Tested Applications:

IHC, FC

Species Specificity:

human

Positive Controls:

IHC : human tonsillitis tissue,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436). It has been reported that PD-1 is heavily glycosylated and migrates with an apparent molecular mass of 47-55 kDa on SDS-PAGE, which is larger than its predicted mass of 32 kDa (PMID: 8671665; 17640856; 17003438).

## Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH 7.3.

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




1x10<sup>6</sup> unstimulated or PHA stimulated human PBMCs were surface stained with 0.25 ug Anti-Human PD-1/CD279 Rabbit Recombinant Antibody (98068-1-RR, Clone:240724G11) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD3. Cells were not fixed.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98068-1-RR (PD-1/CD279 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98068-1-RR (PD-1/CD279 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10<sup>6</sup> PHA stimulated human PBMCs were stained with 0.25 ug Anti-Human PD-1/CD279 Rabbit Recombinant Antibody (98068-1-RR, Clone:240724G11)(left) or 0.25 ug PE Anti-Human PD-1/CD279 (J110) (PE-65119, Clone: J110)(right) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (APC-65569, Clone: OKT3). Cells were not fixed.



1x10<sup>6</sup> unstimulated (black) or PHA stimulated human PBMCs (red) were stained with 0.25 ug Anti-Human PD-1/CD279 Rabbit Recombinant Antibody (98068-1-RR, Clone:240724G11) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). 1x10<sup>6</sup> PHA treated human PBMCs were stained with Isotype Control(blue). Cells were not fixed.



Bilayer interferometry (BLI) kinetic assays of 98068-1-RR against Human PD-1/CD279 were performed. The affinity constant is 0.966 nM.