For Research Use Only

Anti-Human CD79a Rabbit Recombinant **Antibody**

Catalog Number: 98072-1-RR



Purification Method: Protein A purfication

CloneNo.: 241299F3

Recommended Dilutions:

FC (Intra): 0.25 ug per 10^6 cells in a

100 µl suspension

Basic Information

Catalog Number: 98072-1-RR Concentration:

100ug, 1000 ug/ml Source: Rabbit

Isotype:

Calculated MW: 226 aa, 25 kDa

associated alpha

BC113733

GeneID (NCBI):

UNIPROT ID:

Full Name:

P11912

GenBank Accession Number:

CD79a molecule, immunoglobulin-

Applications

Tested Applications: FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): human PBMCs,

Background Information

CD79a, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cells and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells.

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


Selected Validation Data



1x10^6 human PBMCs were intracellularly stained with 0.25 ug Anti-Human CD79a Rabbit RecAb (98072-1-RR, Clone: 241299F3) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with Coralite® Plus 647 Anti-Human CD19. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer. Lymphocytes were gated.



1x10^6 human PBMCs were intracellularly stained with 0.25 ug Anti-Human CD79a Rabbit RecAb (98072-1-RR, Clone: 241299F3) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer. Lymphocytes were gated.



Biolayer interferometry (BLL) kinetic assays of 98072-1-RR against Human CD79a were performed. The affinity constant is 0.151 nM.