

For Research Use Only

Anti-Human SELPLG/CD162 Rabbit Recombinant Antibody

Catalog Number: 98084-1-RR



Basic Information

Catalog Number:

98084-1-RR

Size:

100ug, 1000 μ g/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC029782

GeneID (NCBI):

6404

UNIPROT ID:

Q14242

Full Name:

selectin P ligand

Calculated MW:

412 aa, 43 kDa

Purification Method:

Protein A purification

CloneNo.:

241297F10

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

SELPLG (selectin P ligand), also known as CD162 or PSGL-1, is a 120 kDa type I transmembrane mucin-like glycoprotein. It is an adhesion molecule expressed on the surface of neutrophils, monocytes, and most lymphocytes as a homodimer of approximately 230-250 kDa (PMID: 7545173; 1378449; 9353350). SELPLG plays a critical role in the early stages of inflammation due to its ability to bind E-selectin (CD62E), L-selectin (CD62L), and P-selectin (CD62P) (PMID: 35597982).

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH 7.3.

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human SELPLG/CD162 Rabbit Recombinant Antibody (98084-1-RR, Clone:241297F10), or Isotype Control and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human SELPLG/CD162 Rabbit Recombinant Antibody (98084-1-RR, Clone:241297F10) (red), or Isotype Control (blue) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.



Biolayer interferometry (BLI) kinetic assays of 98084-1-RR against Human SELPLG/CD162 were performed. The affinity constant is 0.699 nM.