For Research Use Only


Anti-Mouse CD40 Rabbit Recombinant Antibody, PBS Only Catalog Number:98122-1-PBS




Basic Information	Catalog Number: 98122-1-PBS	GenBank Accession Number: NM_011611.2	Purification Method: Protein A purfication
	Size: 1mg, 2 mg/ml	GenelD (NCBI): 21939	CloneNo.: 240874G8
	Source: Rabbit	UNIPROT ID: P27512-1	
	lsotype: IgG	Full Name: CD40 antigen	
		Calculated MW: 32 kDa	
Applications	Tested Applications: FC		
	Species Specificity: mouse		
Background Information	CD40, also named as TNFRSF5 and Bp50, is a type I transmembrane protein that belongs to the tumor necrosis factor-R (TNF-R) family (PMID: 10647992). CD40 is expressed on B cells, dendritic cells (DCs), monocytes, platelets, and macrophages as well as by non-hematopoietic cells such as myofibroblasts, fibroblasts, epithelial, and endothelial cells (PMID: 19426221). It is a receptor for TNFSF5/CD40LG. CD40L/CD40 interactions are essential for immunoglobulin (Ig) isotype switching, germinal center formation and the development of B cell memory (PMID: 7516405; 11675497).		
Storage	Storage: Store at -80°C. The product is shipped with i Storage Buffer: PBS Only	ce packs. Upon receipt, store it immediatel	yat-80℃


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD40 Rabbit Recombinant Antibody (98122-1-RR, Clone: 240874G8) or 0.25 ug Isotype Control, and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with APC Anti-Mouse CD19 (1D3) (APC-65290, Clone: 1D3). Cells were not fixed. This data was developed using the same antibody clone with 98122-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 98122-1-RR against Mouse CD40 were performed. The affinity constant is 0.997 nM.

1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD40 Rabbit Recombinant Antibody (98122-1-RR, Clone: 240874G8) (red) or 0.25 ug Isotype Control (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. This data was developed using the same antibody clone with 98122-1-PBS in a different storage buffer formulation.