For Research Use Only

## Rabbit IgG Isotype Control Recombinant proteintech Antibody, PBS Ónly

Catalog Number:98136-1-PBS



**Basic Information** 

Catalog Number: 98136-1-PBS

Concentration:

1mg, 3000  $\,\mu$  g/ml

Source: Rabbit Isotype: GenBank Accession Number:

GeneID (NCBI):

Full Name:

**Purification Method:** Protein A purfication

CloneNo.: 240953C9

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC, FC (Intra), IP, ELISA, ChIP-qPCR

Species Specificity:

n/a

**Background Information** 

This product is an isotype control antibody, which is used to estimate the non-specific binding of target primary antibodies due to Fc receptor binding or other protein-protein interactions. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C  $\,$ 

Storage Buffer: PBS only, pH7.3


## Selected Validation Data




1x10^6 human PBMCs were surface stained with 0.2 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L) (red in histogram), or 0.2 ug Anti-Human CD4 Rabbit Recombinant Antibody (98042-1-RR) and PE-Conjugated Goat Anti-Rabbit IgG(H+L) (green in histogram), or unstained (blue in histogram). Cells were not fixed. This data was developed using the same antibody clone with 98136-1-




IP result of anti-IgG Isotype Control (IP:98136-1-RR, 4ug; Detection:98136-1-RR 1:0) with HepG2 cells lysate 2000 ug. This data was developed using the same antibody clone with 98136-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IgG Isotype Control antibody (98136-1-RR, Clone: 240953C9) at dilution of 1:200 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). NOL11 Recombinant antibody (83391-6-RR) is a positive control. This data was developed using the same antibody clone with 98136-1-PBS in a different storage buffer formulation.




1x10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone:240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.2 ug PGF Recombinant antibody (83170-4-RR, Clone:230532H3) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (green), or unstain (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was



HeLa cells were subjected to SDS PAGE followed by western blot with 98136-1-RR (IgG Isotype Control antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 98136-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 98136-1-RR (IgG Isotype Control antibody) at dilution of 2 ug/mL or 83793-5-RR (Cadherin-17 Recombinant antibody) or unstained (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 98136-1-PBS in a different storage buffer formulation.



Chromatin was prepared from Hela cells.
Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15 µg of cross-linked chromatin, 5 µg of Acetyl-Histone H2A (Lys9) (83041-1-RR) or 5 µg of Normal Rabbit IgG (98136-1-RR), and 20 µl of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real-time PCR. This data was developed using the same antibody clone with 98136-1-PBS in a different storage buffer formulation.