For Research Use Only

Anti-Human CXCR4/CD184 Rabbit Recombinant Antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Protein A purfication

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100 μ l

CloneNo.:

241735F8

suspension

Catalog Number: 98175-1-RR

Basic Information

Catalog Number: 98175-1-RR

Concentration: 100ug, 1000 ug/ml

Source: Rabbit Isotype:

gG cl

Calculated MW: 352 aa, 40 kDa

Applications

Tested Applications:

Species Specificity:

human

GenBank Accession Number:

BC020968 GeneID (NCBI): 7852

UNIPROT ID: P61073

Full Name: chemokine (C-X-C motif) receptor 4

Positive Controls:

FC: human PBMCs,

Background Information

C-X-C chemokine receptor type 4 (CXCR4, also known as CD184) is a widely expressed G protein-coupled seventransmembrane receptor. CXCL12/SDF-1 is the biological ligand for CXCR4. The binding of CXCL12 to CXCR4 induces intracellular signaling through several divergent pathways initiating signals related to chemotaxis, cell survival and/or proliferation, increase in intracellular calcium, and gene transcription (PMID: 20484021). CXCR4 also functions as a coreceptor for HIV-1 entry (PMID: 9427609).

Storage


Storage:

Store at 2 - 8° C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CXCR4/CD184 Rabbit RecAb (98175-1-RR, Clone: 241735F8) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with Coralite® Plus 647 Anti-Human CD19 (4G7) Mouse IgG2a Recombinant Antibody (CL647-65562, Clone: 4G7). Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CXCR4/CD184 Rabbit RecAb (98175-1-RR, Clone: 241735F8) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated.