For Research Use Only

Anti-Human Siglec-7/CD328 Rabbit Recombinant Antibody

Catalog Number: 98183-1-RR



Purification Method:

Protein A purfication

CloneNo.:

241525H1

Basic Information

Catalog Number:

98183-1-RR

Size:

100ug, 1000 ug/ml Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

NM_014385.4 GeneID (NCBI):

27036

UNIPROT ID: Q9Y286-1 Full Name:

sialic acid binding Ig-like lectin 7

Calculated MW:

51kDa

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

Sialic acid binding Ig-like lectin 7 (Siglec-7), also known as CD328 or p75/AIRM-1, is a member of the Siglec family of glycan-recognition proteins. Siglec-7 is a type-I transmembrane protein consisting of three extracellular immunoglobulin-like domains that comprise an N-terminal V-set domain and two C2-set domains, a transmembrane region and a cytoplasmic tail containing two tyrosine residues embodied in immunoreceptor tyrosine-based inhibition motif-like motifs (PMID: 32322597; 10567377). It is mainly expressed on immune cells, with low levels on granulocytes, intermediate levels on monocytes, and relatively high levels on a major subset of natural killer cells and a minor subset of CD8+T cells (PMID: 10567377). Siglec-7 is an inhibitory receptor that negatively regulates the function of NK cells and modulates the immune response through the interaction of sialic acid-containing ligands (PMID: 27312286).

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH 7.3.


Selected Validation Data



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Siglec-7/CD328 Rabbit Recombinant Antibody (98183-1-RR, Clone: 241525H1) or Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with CoraLite® Plus 647 Anti-Human NCAM1/CD56. Cells were not fixed



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Siglec-7/CD328 Rabbit Recombinant Antibody (98183-1-RR, Clone:241525H1) (red) or Rabbit 1gG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue) and PE-conjugated Goat Anti-Rabbit 1gG. Cells were not fixed.



Biolayer interferometry (BLL) kinetic assays of 98183-1-RR against Human Siglec-7/CD328 were performed. The affinity constant is 0.22 nM.