

Anti-Human CD3 Mouse Recombinant Antibody

Catalog Number: 98319-2-MR

Basic Information

Catalog Number:

98319-2-MR

Concentration:

100ug, 1000 ug/ml

Source:

Mouse

Isotype:

IgG2a

GenBank Accession Number:

BC049847

GeneID (NCBI):

916

ENSEMBL Gene ID:

ENSG00000198851

UNIPROT ID:

P07766

Full Name:

CD3e molecule, epsilon (CD3-TCR complex)

Calculated MW:

207 aa, 23 kDa

Purification Method:

Protein A purification

CloneNo.:

SP34

Recommended Dilutions:

IHC: 1:500-1:2000

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

IHC, FC

Species Specificity:

Human, Cynomolgus monkey

Positive Controls:

IHC : human tonsillitis tissue,

FC : human PBMCs,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD3 Mouse RecAb (98319-2-MR, Clone: SP34) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4), and PE-conjugated Goat Anti-Mouse IgG. Cells were then stained with CoraLite® Plus 647 Anti-Human CD19. Cells were not fixed. Lymphocytes were gated.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD3 Mouse RecAb (98319-2-MR, Clone: SP34) (red) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue), and PE-conjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98319-2-MR (CD3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).