Anti-Human KIR2DL3/CD158b2 Rabbit Recombinant Antibody

Catalog Number:98373-3-RR





| Basic Information | Catalog Number: 98373-3-RR | GenBank Accession Number: NM_015868.2 | Purification Method: Protein A purfication |
|------------------------|---|---|---|
| | Concentration: 100ug, 1000 µ g/ml | GenelD (NCBI): 3804 | CloneNo.: 242282E11 |
| | Source: Rabbit | UNIPROT ID: P43628-1 | |
| | Isotype: IgG | Full Name: killer cell immunoglobulin-like receptor, two domains, long cytoplasmic tail, 3 | |
| | | Calculated MW: 38 kDa | |
| Applications | Tested Applications: FC | | |
| | Species Specificity: human | | |
| Background Information | Killer cell immunoglobulin-like receptors (KIRs) are a diverse family of inhibitory and activating receptors expressed on NK cells and a subset of T cells (PMID: 11861603). These polymorphic receptors interact with specific motifs on HLA class I molecules, modulate NK cytolytic activity and are encoded by genes located on chromosome 19q13.4 (PMID: 17069649). KIR2DL3, also known as CD158B2 or NKAT2, is an inhibitory receptor that is specific for HLA-C alleles (HLA-Cw1, HLA-Cw3 and HLA-Cw7) (PMID: 10196125). It is a 341-amino acid transmembrane glycoprotein consisting of an extracellular region containing two C2-type Ig-like domains, a 19-amino acid hydrophobic transmembrane region, and a long cytoplasmic tail with two immunoreceptor tyrosine-based inhibitory motifs (ITIMs). KIR2DL3 inhibits the cytolytic activity of NK cells thus preventing cell lysis (PMID: 7724594). | | |
| Storage | Storage: Store at 2 - 8°C. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide, pH 7.3. | | |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10[^]6 human PBMCs were surface stained with 0.25 ug Anti-Human KIR2DL3/CD158b2 Rabbit RecAb (98373-3-RR, Clone: 242282E11) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with Coralite® Plus 647 Anti-Human NCAM1/CD56. Cells were not fixed. Lymphocytes were gated.

1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human KIR2DL3/CD158b2 Rabbit RecAb (98373-3-RR, Clone: 242282E11) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated.