

For Research Use Only

Anti-Human CSF1R/CD115 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98379-3-PBS



Basic Information

Catalog Number:

98379-3-PBS

Concentration:

1mg, 2 mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC047521

GeneID (NCBI):

1436

UNIPROT ID:

P07333

Full Name:

colony stimulating factor 1 receptor

Calculated MW:

972 aa, 108 kDa

Purification Method:

Protein A purification

CloneNo.:

250001D7

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

The CSF-1 receptor (CSF1R), also called CD115, is activated by the homodimeric growth factors colony-stimulating factor-1 (CSF-1) and interleukin-34 (IL-34). It plays important roles in development and in innate immunity by regulating the development of most tissue macrophages and osteoclasts, of Langerhans cells of the skin, of Paneth cells of the small intestine, and of brain microglia. It also regulates the differentiation of neural progenitor cells and controls functions of oocytes and trophoblastic cells in the female reproductive tract. Owing to this broad tissue expression pattern, it plays a central role in neoplastic, inflammatory, and neurological diseases.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CSF1R/CD115 Rabbit RecAb (98379-3-RR, Clone:250001D7) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98379-3-PBS in a different storage buffer formulation.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CSF1R/CD115 Rabbit RecAb (98379-3-RR, Clone: 250001D7) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98379-3-PBS in a different storage buffer formulation.