

For Research Use Only

Anti-Human IL-15RA Rabbit Recombinant Antibody

Catalog Number: 98380-3-RR



Basic Information

Catalog Number:

98380-3-RR

Concentration:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC107777

GeneID (NCBI):

3601

UNIPROT ID:

Q13261

Full Name:

interleukin 15 receptor, alpha

Calculated MW:

231 aa, 24 kDa

Purification Method:

Protein A purification

CloneNo.:

242689E1

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

FC

Species Specificity:

human

Positive Controls:

FC : LPS and IFN-γ treated human monocytes,

Background Information

Interleukin-15 (IL-15) is a pleiotropic cytokine that plays an important role in both innate and adaptive immunity. IL-15 is mainly produced by activated monocytes, macrophages and dendritic cells, and is structurally similar to IL-2 (PMID: 8178155; 12401478). These two cytokines share the same IL-2/15R β and common γ -chain receptor subunits. In addition, IL-2 and IL-15 have their own private α -chain receptor subunit, IL-2R α and IL-15R α , respectively (PMID: 7641685). The IL-15R α chain is expressed by many cell types including monocytes, DCs, NK, T cells and fibroblasts. Several isoforms of IL-15R α exist and are generated either by alternative splicing or by proteolytic cleavage (PMID: 10480910; 15265897).

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or LPS and IFN- γ treated human monocytes were surface stained with 0.25 ug Anti-Human IL-15RA Rabbit RecAb (98380-3-RR, Clone:242689E1) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD14. Cells were not fixed.



1x10⁶ LPS and IFN- γ treated human PBMCs were surface stained with 0.25 ug Anti-Human IL15RA Rabbit RecAb (98380-3-RR, Clone: 242689E1) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). 1x10⁶ untreated human PBMCs were surface stained with 0.25 ug Anti-Human IL15RA Rabbit RecAb (98380-3-RR, Clone: 242689E1) (black, dashed) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.