

For Research Use Only

Anti-Mouse CXCL4/PF4 Rabbit Recombinant Antibody

Catalog Number: 98397-2-RR



Basic Information

Catalog Number:	98397-2-RR	GenBank Accession Number:	NM_019932	Purification Method:	Protein A purification
Concentration:	1000ug, 1000 μ g/ml	GenID (NCBI):	56744	CloneNo.:	242741F10
Source:	Rabbit	UNIPROT ID:	Q9Z126	Recommended Dilutions:	FC (Intra): 0.25 ug per 10 ⁶ cells in 100 μ l suspension
Isotype:	IgG	Full Name:	platelet factor 4		
Immunogen Catalog Number:	EG3345	Calculated MW:	11kd		

Applications

Tested Applications:	Positive Controls:
FC (Intra)	FC (Intra): Transfected HEK-293T cells,
Species Specificity:	

Background Information

CXCL4, also known as platelet factor 4 (PF-4), is the oldest member of the chemokine family. CXCL4 is a chemokine produced by activated platelets and immune cells involved in pathological conditions such as cancer, infections and inflammatory diseases like systemic sclerosis (SSc), rheumatoid arthritis (RA) and psoriatic arthritis (PsA), among others. CXCL4 is present in the α -granules of platelets in amounts that constitute up to 2% platelet protein mass. CXCL4 plays a determinant role in distinct physiological processes such as in hematopoiesis, angiogenesis, coagulation and modulation of immune responses. For instance, CXCL4 can prevent monocyte apoptosis and promote cell survival, induces the production of TNF and reactive oxygen species (ROS) and promotes monocyte differentiation into a unique macrophage-like phenotype.

Storage


Storage:
Store at 2 - 8°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HEK-293T cells or CXCL4/PF4 transfected HEK-293T cells were intracellularly stained with 0.25 ug Anti-Mouse PF4 Rabbit RecAb (98397-2-RR, Clone: 242741F10) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.

1x10⁶ HEK-293T cells (left) or CXCL4/PF4 transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse PF4 Rabbit RecAb (98397-2-RR, Clone: 242741F10) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.