

For Research Use Only

# Anti-Human PDGFR beta/CD140b Rabbit Recombinant Antibody

Catalog Number: 98405-2-RR



## Basic Information

Catalog Number:

98405-2-RR

Concentration:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG3715

GenBank Accession Number:

BC032224

GeneID (NCBI):

5159

UNIPROT ID:

P09619

Full Name:

platelet-derived growth factor receptor, beta polypeptide

Calculated MW:

1106 aa, 124 kDa

Purification Method:

Protein A purification

CloneNo.:

242711A9

## Applications

Tested Applications:

FC

Species Specificity:

human

## Background Information

PDGFRs are receptor tyrosine kinases with an extracellular ligand-binding domain and a cytoplasmic enzymatic domain. The signaling activity of PDGFRs is tightly regulated by an autoinhibitory allosteric conformation, and requires ligand-induced receptor dimerization at the cell surface to alleviate autoinhibition and achieve physiological activity (PMID: 21664579). PDGFR beta (platelet-derived growth factor receptor-β)/CD140b is a constitutive marker expressed by ASCs in the angiogenic behavior of human retinal endothelial cells (HREs). CD140b (PDGFR β) plays a crucial role in controlling the intrinsic proliferation (PMID: 30976657)

## Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



1x10<sup>6</sup> MG-63 cells were surface stained with 0.25 ug Anti-Human PDGFR beta/CD140b Rabbit RecAb (98405-2-RR, Clone: 242711A9) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.



1x10<sup>6</sup> HEK-293T cells or PDGFR beta/CD140b transfected HEK-293T cells were surface stained with 0.25 ug Anti-Human PDGFR beta/CD140b Rabbit RecAb (98405-2-RR, Clone: 242711A9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.



1x10<sup>6</sup> HEK-293T cells (left) or PDGFR beta/CD140b transfected HEK-293T cells(right) were surface stained with 0.25 ug Anti-Human PDGFRB Rabbit RecAb (98405-2-RR, Clone: 242711A9) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.