For Research Use Only


Anti-Mouse CXCL5 Rabbit Recombinant Α

Antibody, PBS (Catalog Number:98417-2-PR	Dnly		Antibodies ELISA kits Proteins Uni-rAb www.ptglab.com
Basic Information	Catalog Number:	GenBank Accession Number:	Purification Method:
	98417-2-PBS Concentration: 1mg, 2 mg/ml	NM_009141.3 GenelD (NCBI): 20311	Protein A purification CloneNo.: 242737D4
	Source: Rabbit	UNIPROT ID: P50228	
	Isotype: IgG	Full Name: chemokine (C-X-C motif) ligand !	5
	Immunogen Catalog Number: EG1636	Calculated MW: 14KD	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: mouse		
Background Information	Chemokines are a class of pro-inflammatory cytokines that can recruit and activate chemotactic cells. CXCL5 is a member of the chemokine family binding CXCR2, a G-protein coupled receptor. CXCL5 is a chemokine that promotes tumor formation by triggering the migration of immune cells to tumors and promotes immmuno-suppressive characteristics of the tumor microenvironment. CXCL5 can also promote tumor cell metastasis and recruit vascular endothelial cells for angiogenesis.		
Storage	Storage: Store at -80°C. The product is shipped with ice pa	acks. Upon receipt, store it immediatel	ly at −80°C

i ne pro ct is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS only, pH7.3

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 HEK-293T cells (left) or CXCL5-transfected HEK-293T cells (right) with 0.25 ug Anti-Mouse CXCL5 Rabbit RecAb (98417-2-RR, Clone: 242737D4) and APC-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98417-2-PBS in a different storage buffer formulation. 1x10^6 HEK-293T cells (right) or CXCL5transfected HEK-293T cells (left) with 0.25 ug Anti-Mouse CXCL5 Rabbit RecAb (98417-2-RR, Clone: 242737D4) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-2-RR, Clone: 240953A11) (blue), and APC-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98417-2-PBS in a different storage buffer