For Research Use Only

## Anti-Mouse GITR/TNFRSF18 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98466-3-PBS



**Basic Information** 

Catalog Number: 98466-3-PBS Concentration:

1mg, 3 mg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

EG3695

GenBank Accession Number: **Purification Method:** Protein A purification

CloneNo.: 250666B12

Calculated MW: 25kDa

tumor necrosis factor receptor superfamily, member 18

NM\_009400.2

GeneID (NCBI):

**UNIPROT ID:** 

035714-1 Full Name:

21936

**Applications** 

**Tested Applications:** 

Species Specificity:

mouse

## **Background Information**

Glucocorticoid-induced TNFR-related protein (GITR), also known as CD357 or TNFRSF18, is a member of the tumor  $\stackrel{\cdot}{\text{necrosis}} \text{ factor receptor (TNF-R) superfamily. GITR is expressed constitutively at high levels in T regulatory cells} \\$ (Treg cells) and plays a key role in dominant immunological self-tolerance maintained by CD25+CD4+ regulatory T cells (PMID: 11812990). It is expressed at low levels on resting responder T cells. The expression of GITR on T cells can be upregulated upon activation (PMID: 15770698). GITR is activated by GITR ligand (GITRL) which is mainly expressed on APC. GITR-GITRL interactions could co-stimulate both responder T-cell functions and the suppressive functions of Treg cells (PMID: 16868552).

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS only, pH7.3

## **Selected Validation Data**



1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse GITR/TNFRSF18 Rabbit RecAb (98466-3-RR, Clone: 250666B12) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Mouse CD25/IL-2RA (PC61.5) (APC-65137, Clone: PC61.5). CD4+ cells were pregated for analysis. Cells were not fixed. This data was developed using the same antibody



1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse GITR/TNFRSF18 Rabbit RecAb (98466-3-RR, Clone:250666B12) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). CD4+ cells were pre-gated for analysis. Cells were not fixed. This data was developed using the same antibody clone with 98466-3-PBS in a different storage buffer formulation.