

For Research Use Only

Anti-Mouse CD8a Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98472-2-PBS



Basic Information

| | | |
|-------------------------------------|---|--|
| Catalog Number: 98472-2-PBS | GenBank Accession Number: NM_001081110.2 | Purification Method: Protein A purification |
| Concentration: 1mg, 3 mg/ml | GeneID (NCBI): 12525 | CloneNo.: 242879C10 |
| Source: Rabbit | UNIPROT ID: P01731 | |
| Isotype: IgG | Full Name: CD8 antigen, alpha chain | |
| Immunogen Catalog Number: EG3265 | Calculated MW: 27 kDa | |

Applications

Tested Applications:
FC

Species Specificity:
mouse

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C
Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD8a Rabbit RecAb (98472-2-RR, Clone: 242879C10) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with CoraLite® Plus 647 Anti-Mouse CD3. Cells were not fixed. This data was developed using the same antibody clone with 98472-2-PBS in a different storage buffer formulation.



1x10⁶ mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD8a Rabbit RecAb (98472-2-RR, Clone: 242879C10) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. This data was developed using the same antibody clone with 98472-2-PBS in a different storage buffer formulation.