

For Research Use Only

Anti-Mouse CCL8/MCP-2 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98480-1-PBS



Basic Information

Catalog Number:

98480-1-PBS

Concentration:

1mg, 3 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG2771

GenBank Accession Number:

NM_021443.3

GeneID (NCBI):

20307

UNIPROT ID:

Q9Z121

Full Name:

chemokine (C-C motif) ligand 8

Calculated MW:

11 kDa

Purification Method:

Protein A purification

CloneNo.:

250174H6

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Background Information

The cys-cys (C-C) motif chemokine ligand 8 (CCL8, also known as MCP2) is implicated in various inflammatory processes. In bone marrow transplantation, circulating CCL8 levels positively correlate with the severity of graft-versus-host disease.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HEK-293T cells (left) or CCL8/MCP-2-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL8/MCP-2 Rabbit RecAb (98480-1-RR, Clone: 250174H6) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98480-1-PBS in a different storage buffer formulation.

1x10⁶ HEK-293T cells (left) or CCL8/MCP-2-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL8/MCP-2 Rabbit RecAb (98480-1-RR, Clone: 250174H6) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone.