

For Research Use Only

Anti-Human CD55 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98493-3-PBS



Basic Information

Catalog Number: 98493-3-PBS	GenBank Accession Number: BC001288	Purification Method: Protein A purification
Concentration: 1mg, 3 mg/ml	GeneID (NCBI): 1604	CloneNo.: 250682C11
Source: Rabbit	UNIPROT ID: P08174	
Isotype: IgG	Full Name: CD55 molecule, decay accelerating factor for complement (Cromer blood group)	
Immunogen Catalog Number: EG4246	Calculated MW: 41 kDa	

Applications

Tested Applications:
FC

Species Specificity:
human

Background Information

CD55, also known as DAF, is a glycosylphosphatidylinositol (GPI)-anchored surface glycoprotein that is widely distributed on blood, stroma, epithelial, and endothelial cells (PMID: 7517044; 29503741). It can also exist as a soluble form in plasma, urine, saliva, tears, and synovial fluids (PMID: 29503741). CD55 is a complement regulatory protein (PMID: 2469439; 7517044). It inhibits formation of the C3 convertases through binding to C3b and C4b. It also binds the alternate pathway convertase C3bBb, the classical pathway convertase and C4b2a to accelerate their decay (PMID: 17289551). CD55 also serves as a receptor for coxsackieviruses B1, B3, and B5 and several enteroviruses (PMID: 7538177; 7517044).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human blood cells were surface stained with 0.25 ug Anti-Human CD55 Rabbit RecAb (98493-3-RR, Clone: 250682C11) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98493-3-PBS in a different storage buffer formulation.



1x10⁶ human blood cells were surface stained with 0.25 ug Anti-Human CD55 Rabbit RecAb (98493-3-RR, Clone:250682C11) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98493-3-PBS in a different storage buffer formulation.