For Research Use Only

Anti-Human CD73 Rabbit Recombinant Antibody

www.ptglab.com

Catalog Number: 98566-1-RR

Basic Information

Catalog Number: GenBank Accession Number: 98566-1-RR NM_002526.3

GeneID (NCBI): Concentration: 100ug, 1000 $\,\mu$ g/ml

UNIPROT ID: Source: Rabbit P21589-1

Full Name: Isotype:

5'-nucleotidase, ecto (CD73) Calculated MW:

Immunogen Catalog Number:

63 kDa

Applications

Tested Applications:

Species Specificity:

human

Purification Method:

Protein A purification

CloneNo.: 251250H3

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100 μ l

suspension

Positive Controls:

FC: human PBMCs,

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).


Storage


Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH7.3

Selected Validation Data





1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD73 Rabbit RecAb (98566-1-RR, Clone:251250H3) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PEconjugated Goat Anti-Rabbit IgG. Cells were then stained with Coralite® Plus 647 Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (CL647-65569, Clone: OKT3). Cells were not fixed.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD73 Rabbit RecAb (98566-1-RR, Clone: 251250H3) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were not fixed.