

For Research Use Only

Anti-Mouse RON/MST1R Rabbit Recombinant Antibody

Catalog Number: 98587-2-RR



Basic Information

Catalog Number:

98587-2-RR

Concentration:

100ug, 1000 μ g/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG2647

GenBank Accession Number:

NM_009074.2

GeneID (NCBI):

19882

UNIPROT ID:

Q62190-1

Full Name:

macrophage stimulating 1 receptor (c-met-related tyrosine kinase)

Calculated MW:

151 kDa

Purification Method:

Protein A purification

CloneNo.:

250364F1

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100 μ l suspension

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : mouse peritoneal macrophages,

Background Information

RON (also referred to as MST1R, or CD136) is a receptor tyrosine kinase that belongs to the MET proto-oncogene family. The precursor RON is produced as a glycosylated, single chain that is cleaved into a 185-kDa heterodimer of α (35 kDa) and β (150 kDa) subunits connected by a disulfide bond (PMID: 8062829). It is activated upon binding to its ligand, macrophage-stimulating protein (MSP, also known as MST1 and HGFL). RON signalling is essential for embryonic development and regulates many physiological processes including cell survival, migration and differentiation.

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse peritoneal macrophages were surface stained with 0.25 ug Anti-Mouse RON/MST1R Rabbit RecAb (98587-2-RR, Clone: 250364F1) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-conjugated Goat Anti-Rabbit IgG. Cells were co-stained with CoraLite® Plus 750 Anti-Mouse F4/80. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ mouse peritoneal macrophages were surface stained with 0.25 ug Anti-Mouse RON/MST1R Rabbit RecAb (98587-2-RR, Clone: 250364F1) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.