

For Research Use Only

Anti-Human Survivin Rabbit Recombinant Antibody

Catalog Number: 98624-1-RR



Basic Information

Catalog Number: 98624-1-RR	GenBank Accession Number: /	Purification Method: Protein A purification
Concentration: 100ug, 1000 µg/ml	GeneID (NCBI): 332	CloneNo.: 251227G1
Source: Rabbit	UNIPROT ID: O15392-1	Recommended Dilutions: IF/ICC: 1:250-1:1000
Isotype: IgG	Full Name: baculoviral IAP repeat-containing 5	FC (Intra): 0.25 ug per 10 ⁶ cells in a 100 µl suspension
Immunogen Catalog Number: EG4645	Calculated MW: 16kDa	

Applications

Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : MCF-7 cells, A549 cells
Species Specificity: human	FC (Intra) : FC receptor blocked human PBMCs,

Background Information

Survivin, also called BIRC5, is a unique member of the inhibitor of apoptosis (IAP) protein family. Survivin is a 16 kDa anti-apoptotic protein highly expressed during fetal development and cancer cell malignancy, but is completely absent in terminally differentiated cells. The differential expression of survivin in cancer versus normal tissues makes it a useful tool in cancer diagnosis and a promising therapeutic target. Survivin expression is also highly regulated by the cell cycle and is only expressed in the G2-M phase. It is known that survivin localizes to the mitotic spindle by interaction with tubulin during mitosis and may play a contributing role in regulating mitosis. Disruption of survivin-microtubule interactions results in loss of survivin's anti-apoptosis function and increased caspase-3 activity, a mechanism involved in cell death, during mitosis. It also is a direct target gene of the Wnt pathway and is upregulated by beta-catenin.

Storage

Storage:
Store at 2 - 8°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Survivin antibody (98624-1-RR, Clone: 251227G1) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).




1x10⁶ human PBMCs were intracellularly stained with 0.25 ug Anti-Human Survivin Rabbit RecAb (98624-1-RR, Clone:251227G1),or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9)and PE-Conjugated Goat Anti-Rabbit IgG(H+L),Cells were then stained with CoraLite® Plus 647 Anti-Human CD14 (261C) (CL647-65246, Clone: 261C),Cells were fixed and permeabilized with Intracellular Flow Cytometry Fixation & Permeabilization Buffer.



Immunofluorescent analysis of (4% PFA) fixed A549 cells using Survivin antibody (98624-1-RR, Clone: 251227G1) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10⁶ human PBMCs were intracellularly stained with 0.25 ug Anti-Human Survivin Rabbit RecAb (98624-1-RR, Clone:251227G1)(red),or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9)(blue)and PE-Conjugated Goat Anti-Rabbit IgG(H+L).Cells were fixed and permeabilized with Intracellular Flow Cytometry Fixation & Permeabilization Buffer Kit (PF00019).Cells were incubated with FC Receptor Block prior to staining.



Biolayer interferometry (BLI) kinetic assays of 98624-1-RR against Human Survivin were performed. The affinity constant is 3.16 nM.