

For Research Use Only

Anti-Whitlow/218 Linker Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98672-2-PBS



Basic Information

Catalog Number:

98672-2-PBS

Concentration:

1mg, 3 mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Purification Method:

Protein A purification

CloneNo.:

252007G8

Applications

Tested Applications:

WB, FC

Species Specificity:

n/a

Background Information

The 218 Linker (also known as the Whitlow linker) is a synthetic peptide sequence, connecting two protein domains, most commonly the Variable Heavy (VH) and Variable Light (VL) chains of an antibody, to form a stable single-chain variable fragment (scFv) (PMID: 8309948). Its standard amino acid sequence is GSTSGSGKPGSGEGSTKG.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ HEK-293T cells or anti-CD19 (FMC63) CAR with CD28 or CD8a hinge region and transmembrane region transfected HEK-293T cells were surface stained with 0.25 ug Anti-Whitlow/218 Linker Rabbit RecAb (98672-2-RR, Clone:252007G8), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone:240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. This data was developed using the same antibody clone with 98672-2-PBS in a different storage



1x10⁶ HEK-293T cells or anti-CD19 (FMC63) CAR with CD28 or CD8a hinge region and transmembrane region transfected HEK-293T cells were surface stained with 0.25 ug Anti-Whitlow/218 Linker Rabbit RecAb (98672-2-RR, Clone:252007G8) (red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone:240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. This data was developed using the same antibody clone with 98672-2-PBS in a



Various lysates were subjected to SDS PAGE followed by western blot with 98672-2-RR (Whitlow/218 Linker antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 98672-2-PBS in a different storage buffer formulation.