

For Research Use Only

Mouse IgG2b, λ isotype control Recombinant Antibody, PBS Only

Catalog Number: 98739-1-PBS



Basic Information

Catalog Number:

98739-1-PBS

Concentration:

1mg, 3 mg/ml

Source:

Mouse

Isotype:

IgG2a

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Purification Method:

Protein A purification

CloneNo.:

253507A1

Applications

Tested Applications:

FC

Species Specificity:

n/a

Background Information

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Mouse IgG2b, λ isotype control RecAb (98739-1-MR, Clone:253507A1) (red in histogram) and PE-conjugated Goat Anti-Mouse IgG, or 0.25 ug Anti-Human CD4 (OKT4) (65134-1-Ig, Clone: OKT4) (green in histogram), or unstained cells (blue in histogram). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98739-1-PBS in a



1x10⁶ rat splenocytes cells were surface stained with 0.25 ug Mouse IgG2b, λ isotype control RecAb (98739-1-MR, Clone:253507A1) (red in histogram) and PE-conjugated Goat Anti-Mouse IgG, or unstained cells (blue in histogram). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98739-1-PBS in a different storage buffer formulation.



1x10⁶ THP-1 cells were surface stained with 0.25 ug Mouse IgG2b, λ isotype control RecAb (98739-1-MR, Clone:253507A1) (red in histogram) and PE-conjugated Goat Anti-Mouse IgG, or unstained cells (blue in histogram). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98739-1-PBS in a different storage buffer formulation.