

For Research Use Only

Anti-Human IgM Rabbit Recombinant Antibody

Catalog Number: 98744-1-RR



Basic Information

Catalog Number:

98744-1-RR

Concentration:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG6742

GenBank Accession Number:

BC020240

GeneID (NCBI):

3507

Full Name:

immunoglobulin heavy constant mu

Purification Method:

Protein A purification

CloneNo.:

252244H7

Recommended Dilutions:

IHC: 1:250-1:1000

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

IHC, FC

Species Specificity:

human

Positive Controls:

IHC : human tonsillitis tissue,

FC : human PBMCs,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IgM (Immunoglobulin M) is the first immunoglobulin expressed during B cell development. IgM antibodies play an important role in primary immune response involved in early recognition of external invaders like bacteria and viruses, cellular waste and modified self, as well as in recognition and elimination of precancerous and cancerous lesions.

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human Human IgM Rabbit RecAb (98744-1-RR, Clone: 252244H7) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human IgD. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98744-1-RR (IgM antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human Human IgM Rabbit RecAb (98744-1-RR, Clone: 252244H7) (red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98744-1-RR (IgM antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).