

For Research Use Only

Anti-Human uPA/Urokinase Rabbit Recombinant Antibody

Catalog Number: 98824-2-RR



Basic Information

Catalog Number: 98824-2-RR	GenBank Accession Number: NM_002658.6	Purification Method: Protein A purification
Concentration: 100ug, 1000 µg/ml	GeneID (NCBI): 5328	CloneNo.: 252624A9
Source: Rabbit	UNIPROT ID: P00749-1	Recommended Dilutions: FC: 0.25 ug per 10 ⁶ cells in a 100 µl suspension
Isotype: IgG	Full Name: plasminogen activator, urokinase	
Immunogen Catalog Number: EG7083	Calculated MW: 49kDa	

Applications

Tested Applications: FC	Positive Controls: FC : HT-1080 cells,
Species Specificity: human	

Background Information

uPA (Urokinase-type plasminogen activator) is also named as Urokinase or PLAU, and belongs to the peptidase S1 family. uPA initiates a proteolytic cascade, which degrades the extracellular matrix during tissue growth and remodelling. The secreted proform of uPA is activated after binding to a specific high-affinity cell surface receptor. Secreted UPA is exposed to rapid inactivation in the pericellular space by plasminogen activator inhibitor-1 (PAI-1), which is stored in the extracellular matrix bound to vitronectin (PMID: 9665342). It has 2 isoforms produced by alternative splicing. Defects in uPA are the cause of Quebec platelet disorder (QPD).

Storage

Storage:
Store at 2 - 8°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HT-1080 (left) or Ramos cells (right) were surface stained with 0.25 ug Anti-Human uPA/Urokinase Rabbit RecAb (98824-2-RR, Clone: 252624A9) (red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were not fixed.



Biolayer interferometry (BLI) kinetic assays of 98824-2-RR against Human uPA/Urokinase were performed. The affinity constant is below 1 pM.