

APC-conjugated CD206 Polyclonal antibody

Catalog Number: APC-18704

Basic Information

Catalog Number:

APC-18704

Size:

200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002438

GeneID (NCBI):

4360

UNIPROT ID:

P22897

Full Name:

mannose receptor, C type 1

Calculated MW:

166 kDa

Observed MW:

165-180 kDa

Purification Method:

Antigen affinity purification

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Tested Applications:**Species Specificity:**

human, mouse, rat

Background Information

CD206, also named as MMR, CLEC13D and MRC1, is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. CD206 has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. CD206 is a 170 kDa transmembrane glycoprotein which contains 5 domains: an amino-terminal cysteine-rich region, a fibronectin type II repeat, a series of eight tandem lectin-like carbohydrate recognition domains (responsible for the recognition of mannose and fucose), a transmembrane domain, and an intracellular carboxy-terminal tail. It is expressed on most tissue macrophages, in vitro derived dendritic cells, lymphatic and sinusoidal endothelia. This antibody recognizes the intracellular carboxy-terminal part of CD206 and MRC1L1. If protein aggregation exists, for optimal WB detection with this antibody, we recommend adding DTT before boiling the sample to reduce disulfide bonds.

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA.

Selected Validation Data



1×10^6 RAW 264.7 cells were intracellularly stained with 0.4 μ g APC Anti-Human CD206 (APC-18704) (red), or 0.4 μ g Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).