For Research Use Only

APC Anti-Mouse CD107a / LAMP1 (1D4B)

www.ptglab.com

Catalog Number: APC-65050

2 Publications

Basic Information

Catalog Number: APC-65050 Concentration: 100ug, 100 ug/ml

Source: Isotype:

IgG2a, kappa

GenBank Accession Number:

BC006785 GeneID (NCBI): 16783

Full Name:

lysosomal-associated membrane protein 1

UNIPROT ID:

P11438

Purification Method:

Affinity purification

CloneNo.: 1D4B

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications: IF/ICC, FC (Intra) **Cited Applications:**

Species Specificity: mouse Cited Species: mouse

Positive Controls:

IF/ICC: NIH/3T3 cells, FC (Intra): NIH/3T3 cells,

Background Information

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).

Notable Publications

Author	Pubmed ID	Journal	Application
Kai Li	39952937	Nat Commun	IF
Lei Wang	37979663	Pharmacol Res	IF


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using APC-65050 (CD107a antibody) at dilution of 1:100.



1x10^6 NIH/3T3 cells were intracellularly stained with 0.4 ug APC Anti-Mouse CD107a / LAMP1 (APC-65050, Clone:1D4B) (red) or 0.4 ug APC Rat IgG2a Isotype Control (2A3) (APC-65209, Clone: 2A3) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).