

For Research Use Only

APC Anti-Mouse TIGIT (1G9)

Catalog Number: APC-65079



Basic Information

Catalog Number:

APC-65079

Concentration:

100ug, 0.2 mg/ml

Source:

Mouse

Isotype:

IgG1, kappa

GenBank Accession Number:

NM_001146325

GeneID (NCBI):

100043314

UNIPROT ID:

P86176

Full Name:

T cell immunoreceptor with Ig and ITIM domains

Purification Method:

Affinity purification

CloneNo.:

1G9

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in 100 µl suspension

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : plate-bound Anti-CD3 and Anti-CD28 stimulated BALB/c mouse splenocytes,

Background Information

TIGIT (T-cell immunoreceptor with Ig and ITIM domains), also known as VSIG9 or VSTM3, is an immune receptor expressed on T cells, including Treg and memory subsets, as well as on NK cells (PMID: 19011627). It contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT binds to poliovirus receptor (PVR, also called CD155) with high affinity, and also to PVRL2 (CD112) with lower affinity. The interaction of TIGIT with PVR on dendritic cells increases the secretion of IL-10 and decreases the secretion of proinflammatory cytokine and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells. TIGIT can inhibit NK cytotoxicity directly through its ITIM (PMID: 19815499).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.1% gelatin, pH7.2

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ unstimulated or plate-bound Anti-CD3 and Anti-CD28 stimulated BALB/c mouse splenocytes were surface stained with 0.25 ug CoraLite@488 Anti-Mouse CD3 (CL488-65077, Clone: 17A2) and 0.25 ug APC Anti-Mouse TIGIT (APC-65079, Clone: 1G9). Cells were not fixed.