For Research Use Only

APC Anti-Human CD8a (Hit8a)

Catalog Number: APC-65113



Basic Information

Catalog Number:

APC-65113

Size:

100tests, 5 μ l/test

Source: Mouse Isotype:

IgG1, kappa

ENSG00000153563 UNIPROT ID: P01732

BC025715

GeneID (NCBI):

ENSEMBL Gene ID:

CD8a molecule Calculated MW: 235 aa, 26 kDa

Full Name:

GenBank Accession Number:

Purification Method:

Affinity purification

CloneNo.: Hit8a

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD3 and 5 ul APC Anti-Human CD8a (APC-65113, Clone:Hit8a) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul APC Anti-Human CD8a (APC-65113, Clone:Hit8a) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.