For Research Use Only

APC Anti-Mouse CD134/OX40 (OX-86)

Catalog Number: APC-65136 1 Publications




| Basic Information | Catalog Number: GenBank Accession Number: APC-65136 BC065782 | | cession Number: | Purification Method: Affinity purification | |
|------------------------|---|-----------|-----------------|---|--|
| | Size: GeneID (NCBI): | | 31): | CloneNo.: | |
| | 100ug, 0.2 mg/ml | | | OX-86 | |
| | Source: UNIPROT ID: | | | Excitation/Emission maxima wavelengths: 650 nm / 660 nm | |
| | Rat | P47741 | | | |
| | Isotype: Full Name: IgG1, kappa tumor necrosis factor receptor superfamily, member 4 | | • | | |
| Applications | Tested Applications: FC | | | | |
| | Cited Applications: FC | | | | |
| | Species Specificity: mouse | | | | |
| | Cited Species: | | | | |
| | mouse | | | | |
| Background Information | CD134, also known as OX40 and TNFRSF4, is a member of the TNFR-superfamily of receptors (PMID: 2828930; 9766631). It is a type I transmembrane protein predominantly expressed on activated T cells which include CD4 and CD8 T cells, Th2, Th1, and Th17 cells, as well as regulatory T cells (Tregs) (PMID: 20307208). CD134 is activated by its cognate ligand CD134L (OX40L) and functions as a T cell co-stimulatory molecule (PMID: 26215166). CD134- CD134L interactions have been proposed as a potential therapeutic target for treating autoimmune diseases, cancer and infectious disease (PMID: 26215166; 19426222). | | | | |
| Notable Publications | Author | Pubmed ID | Journal | Application | |
| | Yui Maehara | 38839354 | Biomed Res | FC | |
| Storage | Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.1% sodium azide and 0.5% BSA. | | | | |

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 unstimulated or anti-CD3/CD28 stimulated (2 days) mouse splenocytes were surface costained with FITC Plus Anti-Mouse CD3 (17A2) (FITC-65077, Clone: 17A2) and 0.2 ug APC Anti-Mouse CD134 (APC-65136, Clone: 0X-86). Cells were not fixed.