

For Research Use Only

APC Anti-Mouse PD-1/CD279 (RMP1-30)



Catalog Number: APC-65142

1 Publications

Basic Information

Catalog Number: APC-65142	GenBank Accession Number: BC119179	Purification Method: Affinity purification
Size: 100ug, 0.2 mg/ml	GeneID (NCBI): 18566	CloneNo.: RMP1-30
Source: Rat	UNIPROT ID: Q02242	Excitation/Emission maxima wavelengths: 650 nm / 660 nm
Isotype: IgG2b, kappa	Full Name: programmed cell death 1	

Applications

Tested Applications:
FC

Cited Applications:
FC

Species Specificity:
Mouse

Cited Species:
mouse

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Notable Publications

Author	Pubmed ID	Journal	Application
Qing Zhang	35765095	Stem Cell Res Ther	FC

Storage


Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
Phosphate based buffer with 0.09% sodium azide and 0.1% gelatin, pH 7.2.

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ unstimulated or anti-CD3/CD28 treated (2 days) BALB/C mouse splenocytes were surface stained with 0.2 ug APC Anti-Mouse PD-1/CD279 (APC-65142, Clone: RMP1-30), and 0.5 ug FITC Plus Anti-Mouse CD3 (17A2) (FITC-65077, Clone: 17A2). Cells were not fixed.