For Research Use Only

APC Anti-Human PD-1/CD279 (EH12.2H7)



Catalog Number: APC-65168

Basic Information

Catalog Number:

APC-65168

Size:

100tests, 10 μ l/test

Source: Mouse Isotype:

lgG1, kappa

GenBank Accession Number:

BC074740 GeneID (NCBI):

5133

UNIPROT ID: Q15116 Full Name:

programmed cell death 1

Calculated MW: 288 aa, 32 kDa

Purification Method:

The purified antibody is conjugated with allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion

chromatography.

CloneNo.: EH12.2H7

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications:

FC

Species Specificity:

Non-human primates, Human

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide.

Selected Validation Data



1X10^6 human peripheral blood mononuclear cells (PBMCs) were surface stained with FITC Anti-Human CD3 (FITC-65151, clone UCHT1) and 10.00 ul/test APC Anti-Human PD-1/CD279 (APC-65168, Clone:EH12.2H7). Cells were not fixed.



1X10^6 human peripheral blood mononuclear cells (PBMCs) were surface stained with FITC Anti-Human CD3 (FITC-65151, clone UCHT1) and APC mouse IgG1 isotype control antibody. Cells were not fixed.