## For Research Use Only

## APC Anti-Human CD64 (10.1)

Catalog Number: APC-65253



**Purification Method:** 

Excitation/Emission maxima

CloneNo.:

wavelengths: 650 nm / 660 nm

10.1

**Basic Information** 

Catalog Number: GenBank Accession Number:

APC-65253 BC032634

Concentration: GeneID (NCBI):
100tests, 5 ul/test 2209

 Source:
 ENSEMBL Gene ID:

 Mouse
 ENSG00000150337

 Isotype:
 UNIPROT ID:

 IgG1, kappa
 P12314

Full Name:

Fc fragment of IgG, high affinity Ia,

receptor (CD64)
Calculated MW:
374 aa, 43 kDa

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

human

## **Background Information**

Fc  $\gamma$  receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fc  $\gamma$  RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 5 ul APC Anti-Human CD64 (APC-65253, Clone:10.1) or APC Mouse IgG1 Isotype Control (APC-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10^6 human PBMCs were surface stained with CL488 Anti-Human CD11b (CL488-65166) and 5 ul APC Anti-Human CD64 (APC-65253, Clone:10.1) or APC Mouse IgG1 Isotype Control (APC-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.