For Research Use Only

FcZero-rAb™ APC Anti-Human CD11c (3.9) Rabbit IgG Recombinant Antibody

Catalog Number: APC-FcA65568



Basic Information

Catalog Number: APC-FcA65568

Concentration: 100tests, 5 ul/test

Rabbit Isotype:

IgG

Source:

GenBank Accession Number:

BC038237 GeneID (NCBI): 3687

ENSEMBL Gene ID: ENSG00000140678 Full Name:

integrin, alpha X (complement component 3 receptor 4 subunit)

Calculated MW: 1169 aa, 129 kDa Purification Method:

Protein A purification

CloneNo.:

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11c, also known as integrin α X, is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α / β heterodimer with CD18 (integrin β 2). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 human PBMCs were surface stained with PE Anti-Human CD14, and 5 ul APC Anti-Human CD11c (3.9) Rabbit IgG RecAb (APC-FcA65568, Clone: 3.9) or APC Rabbit IgG Isotype Control RecAb (APC-FcA98136, Clone: 240953C9). Cells were not fixed.