

For Research Use Only

FcZero-rAb™ APC Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG Recombinant Antibody

Catalog Number: APC-FcA65663



Basic Information

Catalog Number:

APC-FcA65663

Concentration:

100ug, 200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC031711

GeneID (NCBI):

14960

UNIPROT ID:

P14434

Full Name:

histocompatibility 2, class II antigen A, alpha

Purification Method:

Protein A purification

CloneNo.:

M5/114.15.2

Recommended Dilutions:

FC: 0.1 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : mouse splenocytes,

Background Information

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 antibody is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse splenocytes were surface stained with FcZero-rAb™ PE Anti-Mouse CD3e (145-2C11), and 0.1 ug APC Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (APC-FcA65663, Clone: M5/114.15.2) or FcZero-rAb™ APC Rabbit IgG Isotype Control Recombinant Antibody (APC-FcA98136, Clone: 240953C9). Cells were not fixed.



1x10⁶ mouse splenocytes were surface stained with 0.1 ug APC Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (APC-FcA65663, Clone: M5/114.15.2) (red) or FcZero-rAb™ APC Rabbit IgG Isotype Control Recombinant Antibody (APC-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.