For Research Use Only

FcZero-rAb™ APC-Cyanine7 Anti-Mouse CD206 Rabbit Recombinant Antibody

Catalog Number: AY7-FcA98031



Basic Information

Catalog Number: GenBank Accession Number: AY7-FcA98031 NM_008625.2

GeneID (NCBI): Concentration: CloneNo.: 100ug, 200 ug/ml 17533 240344D12

UNIPROT ID: Source: Recommended Dilutions:

Rabbit Q2HZ94 FC: 0.25 ug per 10^6 cells in a 100 μ l suspension

Isotype: Full Name:

mannose receptor, C type 1 Excitation/Emission maxima

wavelengths: Calculated MW: Immunogen Catalog Number: 650 nm / 778 nm EG0790 165 kDa

Applications

Tested Applications:

Species Specificity:

mouse

Positive Controls:

FC: Balb/c mouse peritoneal macrophages,

Purification Method:

Protein A purification

Background Information

CD206, also named as MMR, CLEC13D and MRC1, is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. CD206 has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. CD206 is a 170 kDa transmembrane protein which contains 5 domains: an amino-terminal cysteine-rich region, a fibronectin type II repeat, a series of eight tandem lectin-like carbohydrate recognition domains (responsible for the recognition of mannose and fucose), a transmembrane domain, and an intracellular carboxy-terminal tail. It is expressed on most tissue macrophages, in vitro derived dendritic cells, lymphatic and sinusoidal endothelia.


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data





1x10^6 Balb/c mouse peritoneal macrophages were surface stained with 0.25 ug APC-Cyanine7 Anti-Mouse CD206 Rabbit RecAb (AY7-FcA98031, Clone: 240344D12) or 0.25 ug FcZero-rAb™ APC-Cyanine7 Rabbit IgG Isotype Control Recombinant Antibody (AY7-FcA98136, Clone: 240953C9). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed.



1x10^6 Balb/c mouse peritoneal macrophages were surface stained with 0.25 ug APC-Cyanine7 Anti-Mouse CD206 Rabbit RecAb (AY7-FcA98031, Clone: 240344D12) (red) or 0.25 ug FcZero-rAb™ APC-Cyanine7 Rabbit IgG Isotype Control Recombinant Antibody (AY7-FcA98136, Clone: 240953C9) (blue). Cells were incubated with human Fc blocking Reagent prior to staining. Cells were not fixed.