FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human TMC4 protein



297 g able lyophilized es: n ssion Source: -derived, PGEX-4T, with N-terminal GST. gical Activity: ested toxin Level: e contact the lab for more information estitution: estitute at 0.25 μg/ μ l in 200 μ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect basequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the based of the final solution will vary according to blume used).	
able lyophilized es: n ssion Source: -derived, PGEX-4T, with N-terminal GST. gical Activity: ested toxin Level: e contact the lab for more information stitution: ustitute at 0.25 µg/ µ l in 200 µ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	QTELQDPHPSRELPWPMQARRAHRQRNASRDQVVY GSGTKTDRWARLLRRSKEKTKEGLRS (1-135 aa encoded by BC025323) Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
able lyophilized es: n ssion Source: -derived, PGEX-4T, with N-terminal GST. gical Activity: ested toxin Level: e contact the lab for more information stitution: ustitute at 0.25 µg/ µ l in 200 µ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	(1-135 aa encoded by BC025323) Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
es: n ssion Source: derived, PGEX-4T, with N-terminal GST. gical Activity: ested toxin Level: e contact the lab for more information stitution: stitute at 0.25 µg/ µ l in 200 µ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
n ssion Source: derived, PGEX-4T, with N-terminal GST. gical Activity: ested oxin Level: e contact the lab for more information astitution: stitute at 0.25 µg/ µ l in 200 µ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect basequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
a-derived, PGEX-4T, with N-terminal GST. gical Activity: ested oxin Level: e contact the lab for more information estitution: estitute at 0.25 μg/ μ l in 200 μ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect basequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
sted oxin Level: e contact the lab for more information stitution: stitute at 0.25 μg/ μ l in 200 μ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the poncentration of the final solution will vary according to	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
e contact the lab for more information stitution: Institute at 0.25 µg/ µ l in 200 µ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effect bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the poncentration of the final solution will vary according to	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
stitute at 0.25 μg/ μ l in 200 μ l sterile water for short- storage. reconstitution with sterile water, if glycerol has no effec bsequent experiments, it is recommended to add an volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
volume of glycerol for long-term storage (see Stability torage for more details). fferent concentration is needed for your purposes please t the reconstitution volume as required (please note: the uncentration of the final solution will vary according to	
Centrifuge vial before opening. When reconstituting,	
y pipet and wash down the sides of the vial to ensure ful ery of the protein into solution.	l
by SDS-PAGE with Coomassie Brilliant Blue staining.	
urified protein was Lyophilized from sterile PBS (58mM l lose and 5 % mannitol are added as protectant before ly	•
for up to 12 months at -20°C to -80°C as lyophilized pow	der.
at 2-8°C for (1-2 weeks). Term Storage:	
74 kDa→	
66 kDa→	
i	Term Storage: at 2-8°C for (1-2 weeks). Term Storage: ot and store at -20°C to -80°C for up to 3 months, buffer ining 50% glycerol is recommended for reconstitution. d repeat freeze-thaw cycles.

20 kD

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.