FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human GRIK5 protein



Basic Information	Catalog Number:	Peptide Sequence:
	Form: Available lyophilized	d MRLSNGKLYSAGAGGDAGSAGGPQRLLDDPGPPSGA RPAAPTPCTHVRVCQECRRIQALRASGAGAPPRGLG VPAEATSPPRPRPGPAGPRELAEHE (845-980 aa encoded by NM_001301030)
	Available lyophilized Species:	
	human	
	Expression Source: <i>e coli-</i> -derived, PGEX-4T, with N-terminal GST.	
	Biological Activity: Not tested	
	Endotoxin Level: Please contact the lab for more information	
Reconstitution and Storage	Reconstitution: Reconstitute at 0.25 μg/ μl in 200 μl sterile water for short- term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.	Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
Purity	85%, by SDS-PAGE with Coomassie Brilliant Blue staining.	
Formulation	The purified protein was Lyophilized from sterile PBS (58mM Na2HPO 4, 17mM NaH2PO 4, 68mM NaCl, pH8.). 5 % trehalose and 5 % mannitol are added as protectant before lyophilization. The elution buffer contain 100mM GSH.	
Stability and Storage	Store for up to 12 months at -20°C to -80°C as lyophilized powder.	
Storage of Reconstituted Protein	Short Term Storage: Store at 2-8°C for (1-2 weeks). Long Term Storage: Aliquot and store at -20°C to -80°C for up to 3 months, buffer containing 50% glycerol is recommended for reconstitution. Avoid repeat freeze-thaw cycles.	
Selected Validation Data	74 kDa→ 43 kDa→ ←41 kDa	

28 kDa-

20 kDa-

14 kDa