FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human SMS protein



Basic Information	Catalog Number:	Peptide Sequence:			
	Ag8767	MAAARHSTLDFMLGAKADGETILKGLQSIFQEQGMA ESVHTWQDHGYLATYTNKNGSFANLRIYPHGLVLLDL			
	Size: 50 μg	QSYDGDAQGKEEIDSILNKVEERMKELSQDSTGRVKR			
	Solveg LPPIVRGGAIDRYWPTADGRLVEYDIDEVVYD Form: QNIKILHSKQFGNILILSGDVNLAESDLAYTRA Available lyophilized EDYTGKDVLILGGGDGGILCEIVKLKPKMVTM Species: MVIDGCKKYMRKTCGDVLDNLKGDCYQVLIE human KRYAKEGREFDYVINDLTAVPISTSPEEDSTW	LPPIVRGGAIDRYWPTADGRLVEYDIDEVVYDEDSPY			
		MVIDGCKKYMRKTCGDVLDNLKGDCYQVLIEDCIPVL KRYAKEGREFDYVINDLTAVPISTSPEEDSTWEFLRLIL DLSMKVLKQDGKYFTQGNCVNLTEALSLYEEQLGRLY CPVEFSKEIVCVPSYLELWVFYTVWKKAKP			
			Endotoxin Level: Please contact the lab for more information		
			Reconstitution and Storage	Reconstitution: Reconstitute at 0.25 μg/μl in 200 μl sterile water for short- term storage.	Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the
				After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details).	recommended temperature (see below).
	If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used).				
Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.					
Purity	85%, by SDS-PAGE with Coomassie Brilliant Blue staining.				
Formulation	The purified protein was Lyophilized from sterile PBS (58mM Na2 trehalose and 5 % mannitol are added as protectant before lyophi	•			
Stability and Storage	Store for up to 12 months at -20°C to -80°C as lyophilized powder.				
Storage of	Short Term Storage: Store at 2-8°C for (1-2 weeks).				
Reconstituted Protein	Aliquot and store at -20°C to -80°C for up to 3 months, buffer containing 50% glycerol is recommended for reconstitution. Avoid repeat freeze-thaw cycles.				
Selected Validation Data					
	74 kDa→ 66 kDa→ ←65 kDa				
	$\begin{array}{c} 43 \text{ kDa} \rightarrow \\ 28 \text{ kDa} \rightarrow \end{array}$				

20 kDa