## Recombinant human CXXC5 protein



**Basic Information** 

Catalog Number:

Ag9949

Size:

50 μg Form:

Available lyophilized

Species:

human

**Expression Source:** 

e coli.-derived, PET28a, with N-terminal 6\*His.

Biological Activity:

Not tested

Endotoxin Level:

Please contact the lab for more information

Peptide Sequence:

MMGGESADKATAAAAAASLLANGHDLAAAMAVDKS
NPTSKHKSGAVASLLSKAERATELAAEGQLTLQQFAQ
STEMLKRVVQEHLPLMSEAGAGLPDMEAVAGAEALN
GQSDFPYLGAFPINPGLFIMTPAGVFLAESALHMAGL
AEYPMQGELASAISSGKKKRKRCGMCAPCRRRINCEQ
CSSCRNRKTGHQICKFRKCEELKKKPSAALEKVMLPTG
AAFRWFO

(1-227 aa encoded by BC017439)

Reconstitution and Storage

## Reconstitution:

Reconstitute at 0.25 µg/  $\mu$  l in 200  $\,\mu$  l sterile water for short-term storage.

After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details).

If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used).

Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).

Purity

85%, by SDS-PAGE with Coomassie Brilliant Blue staining.

**Formulation** 

The purified protein was Lyophilized from sterile PBS (58mM Na2HPO 4,17mM NaH2PO 4, 68mM NaCl, pH7.4). 5 % trehalose and 5 % mannitol are added as protectant before lyophilization. The elution buffer contain 300mM imidazole.

Stability and Storage

Store for up to 12 months at -20°C to -80°C as lyophilized powder.

Storage of Reconstituted Protein

Short Term Storage:

Store at 2-8°C for (1-2 weeks).

Long Term Storage:

Aliquot and store at -20°C to -80°C for up to 3 months, buffer containing 50% glycerol is recommended for reconstitution. Avoid repeat freeze-thaw cycles.

Selected Validation Data

