For Research Use Only

Biotin-conjugated CXCR4 Polyclonal antibody



Purification Method:

IHC 1:50-1:500

Positive Controls:

IHC: mouse spleen tissue,

Antigen affinity purification

Recommended Dilutions:

Catalog Number: Biotin-11073

Basic Information

Catalog Number: GenBank Accession Number: Biotin-11073 BC020968

 Biotin-11073
 BC020968

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 7852

Source: UNIPROT ID: Rabbit P61073
Isotype: Full Name:

IgG chemokine (C-X-C motif) receptor 4

Immunogen Catalog Number:Calculated MW:AG1528352 aa, 40 kDaObserved MW:

Observed M 60-70 kDa

Applications

Tested Applications:

IHC

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

C-X-C chemokine receptor type 4 (CXCR4) is a widely expressed G protein-coupled seven-transmembrane receptor. CXCL12/SDF-1 is the biological ligand for CXCR4. The binding of CXCL12 to CXCR4 induces intracellular signaling through several divergent pathways initiating signals related to chemotaxis, cell survival and/or proliferation, increase in intracellular calcium, and gene transcription (PMID: 20484021). CXCR4 also functions as a coreceptor for HIV-1 entry (PMID: 9427609). CXCR4 has a calculated molecular weight of 40 kDa. In addition to the predicted species of 45-47 kDa for glycosylated CXCR4 monomers, due to ubiquitination, dimeration, and different degrees of glycosylation, additional species can also exist and have been reported in some researches, including 67 kDa (PMID: 23917520), 55, 67, 87 kDa (PMID: 20028517), 80 kDa (PMID: 10506573), 47, 50, 62, and 98 kDa (PMID: 16204649).

Storage

Storage:


Store at -20°C. Avoid exposure to light.


Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data







Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using Biotin-11073 (CXCR4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).