For Research Use Only

Biotin Anti-Human MICA/MICB (6D4)

Catalog Number:Biotin-65161



Basic Information	Catalog Number: Biotin-65161	GenBank Accession Number: BC016929	Purification Method: The purified antibody is conjugated
	Concentration: 100ug, 1 mg/ml Source: Mouse	GenelD (NCBI): 100507436 UNIPROT ID: Q29983	with biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin. CloneNo.: 6D4
	Calculated MW: 383 aa, 43 kDa		
	Applications	Tested Applications: IHC, FC	
Species Specificity: human		IHC : human tonsillitis tissue, human colon tissue FC : HeLa cells,	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	Human MHC class I chain-related genes located within the HLA class I region of chromosome 6 encode MHC class I chain-related A and B (MICA and MICB) (PMID: 11429322). MICA and MICB are stress-inducible surface molecules that are not associated with β 2-microglobulin and do not present peptides (PMID: 9497295). They are expressed in intestinal epithelium and many epithelial tumors (PMID: 10359807). MICA and MICB are ligands for NKG2D which is an activating receptor that is expressed on most natural killer (NK) cells, CD8 α β T cells, and γ δ T cells (PMID: 10426993; 11491531). The monoclonal antibody 6D4 recognizes MICA and MICB.		
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide, pH7.3		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using Biotin-65161 (MICA/MICB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using Biotin-65161 (MICA/MICB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

