For Research Use Only

Biotin Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody

Catalog Number:Biotin-65559



Basic Information

Catalog Number: Biotin-65559

Concentration:

100ug, 500 µg/ml Source:

Mouse
Isotype:
IgG2a

GenBank Accession Number: BC025715

GeneID (NCBI):

925

P01732

ENSEMBL Gene ID: ENSG00000153563 UNIPROT ID:

Full Name: CD8a molecule Calculated MW: 235 aa, 26 kDa Purification Method: Protein A purification

CloneNo.: UCHT4

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 5 ul Biotin Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody (Biotin-65559, Clone:UCHT4) and Coralite@488-conjugated streptavidin, or unstained cells. Cells were costained with APC Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 5 ul Biotin Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody (Biotin-65559, Clone: UCHT4) and CoraLite® 488-conjugated streptavidin (red), or unstained cells (blue). Cells were not fixed. Lymphocytes were gated.