

For Research Use Only

Biotin-conjugated IgD Monoclonal antibody



Catalog Number: Biotin-67538

Basic Information

Catalog Number: Biotin-67538	GenBank Accession Number: BC021276	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 3495	CloneNo.: 1D1B12
Source: Mouse	Full Name: immunoglobulin heavy constant delta	Recommended Dilutions: IHC 1:50-1:500
Isotype: IgG2a	Calculated MW: 573 aa, 63 kDa	
Immunogen Catalog Number: AG9614	Observed MW: 55-70 kDa	

Applications

Tested Applications:

IHC

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC : human tonsillitis tissue,

Background Information

IgD is the major antigen receptor isotype on the surface of most peripheral B-cells. The function of IgD is to signal the B cells to be activated. The relative molecular mass and half-life of secreted IgD is 185 kDa and 2.8 days, respectively. Secreted IgD is produced as a monomeric antibody with two heavy chains of the delta (δ) class, and two Ig light chains. This antibody detects the heavy chain (55-70 kDa) of IgD and secreted IgD (185 kDa) .

Storage

Storage:

Store at -20°C. Avoid exposure to light.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using Biotin-67538 (human IgD antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).