

For Research Use Only

FcZero-rAb™ Biotin Anti-Mouse CD80 Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98158



Basic Information

Catalog Number:

Biotin-FcA98158

Concentration:

100ug, 500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG0840

GenBank Accession Number:

NM_009855.2

GeneID (NCBI):

12519

Full Name:

CD80 antigen

Calculated MW:

35 kDa

Purification Method:

Protein A purification

CloneNo.:

241371E12

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : mouse peritoneal macrophages,

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse peritoneal macrophages were surface stained with 0.25 ug Biotin Anti-Mouse CD80 Rabbit RecAb (Biotin-FcA98158, Clone: 241371E12) or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9), and PE-conjugated streptavidin. Cells were not fixed.



1x10⁶ mouse peritoneal macrophages were surface stained with 0.25 ug Biotin Anti-Mouse CD80 Rabbit RecAb (Biotin-FcA98158, Clone:241371E12)(red) or 0.25 ug Rabbit IgG Isotype Control RecAb (Fc98136-1-RR, Clone: 240953C9) (blue), and PE-conjugated streptavidin. Cells were not fixed.