

For Research Use Only

FcZero-rAb® Biotin Anti-Human CD16a Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98293



Basic Information

Catalog Number:

Biotin-FcA98293

Concentration:

100ug, 500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG2488

GenBank Accession Number:

GeneID (NCBI):

2214

ENSEMBL Gene ID:

ENSG00000203747

Full Name:

Fc fragment of IgG, low affinity IIIa, receptor (CD16a)

Calculated MW:

29 kDa

Purification Method:

Protein A purification

CloneNo.:

242426G11

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

-

Applications

Tested Applications:

FC, ELISA

Species Specificity:

human

Positive Controls:

FC : human peripheral blood leukocytes,

Background Information

CD16 is a 50-70-kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD16 has been identified as Fc receptors FcγRIIIa (CD16a) and FcγRIIIb (CD16b), encoded by two nearly identical genes, FCGR3A and FCGR3B. This antibody is specific for CD16a and does not recognize CD16b.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Biotin Anti-Human CD16a Rabbit RecAb (Biotin-FcA98293, Clone: 242426G11) or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9). Cells were not fixed.



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Biotin Anti-Human CD16a Rabbit RecAb (Biotin-FcA98293, Clone: 242426G11) (red) or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.