For Research Use Only

CoraLite® Plus 405 Anti-Human CD81 (5A6) Mouse IgG2a Recombinant Antibody





Catalog Number:CL405-65644

Basic Information	Catalog Number: CL405-65644	GenBank Accession Number: BC002978	Purification Method: Protein A purification
	Size: 100tests, 5 ul/test	GeneID (NCBI): 975	CloneNo.: 5A6
	Source: Mouse	ENSEMBL Gene ID: ENSG00000110651	Excitation/Emission maxima wavelengths: 399 nm / 422 nm
	Isotype: IgG2a	Full Name: CD81 molecule Calculated MW: 26 kDa	
Species Specificity: human			
Background Information	CD81 (also known as TAPA1 or TSPAN28) is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Many of these members are expressed on leukocytes and have been implicated in signal transduction, cell-cell interactions, and cellular activation and development. CD81 is involved in signal transduction and cell adhesion in the immune system (PMID: 9597125). CD81 has also been identified as en essential receptor for HCV (hepatitis C virus) (PMID: 21428934).		
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide and 0.5% BSA.		

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10[^]6 human peripheral blood leukocytes were surface stained with 5 ul Coralite® Plus 405 Anti-Human CD81 (5A6) Mouse IgG2a RecAb (CL405-65644, Clone: 5A6), or Coralite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.

1x10^6 human peripheral blood leukocytes were surface stained with 5 ul CoraLite® Plus 405 Anti-Human CD81 (5A6) Mouse IgG2a RecAb (CL405-65644, Clone: 5A6) (red), or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.