For Research Use Only

CoraLite® Plus 405 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a Recombinant Antibody



Catalog Number: CL405-65663

Basic Information

Catalog Number:GenBank Accession Number:Purification Method:CL405-65663BC031711Protein A purification

 Concentration:
 GeneID (NCBI):
 CloneNo.:

 100ug, 500 ug/ml
 14960
 M5/114.15.2

Source:UNIPROT ID:Recommended Dilutions:RatP14434FC: 0.25 ug per 10^6 cells in a 100 μl

Isotype: Full Name: suspension

IgG2a histocompatibility 2, class II antigen Excitation/Emission maxima

A, alpha wavelengths: 399 nm / 422 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

Background Information

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 antibody is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 mouse splenocytes were surface stained with CoraLite® Plus 488 Anti-Mouse CD3, and 0.25 ug CoraLite® Plus 405 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a RecAb (CL405-65663, Clone:M5/114.15.2), or CoraLite® Plus 405 Rat IgG2a Isotype Control (2A3) (CL405-65209, Clone: 2A3). Cells were not fixed.



CL405-65663 MHC Class II (I-A/I-E),PB450-A

1x10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 405 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a RecAb (CL405-6566, Clone:M5/114.15.2) (red) or CoraLite® Plus 405 Rat IgG2a Isotype Control (2A3) (CL405-65209, Clone: 2A3) (blue). Cells were not fixed.