## For Research Use Only

## CoraLite® Plus 488-conjugated HO-1/HMOX1 Polyclonal antibody


Catalog Number:CL488-10701 Featured Product


| proteintech®                       |
|------------------------------------|
| Antibodies   ELISA kits   Proteins |

| Basic Information     | Catalog Number:<br>CL488-10701   | GenBank Accession Number:<br>BC001491                   | Purification Method:<br>Antigen affinity purification |  |
|-----------------------|--|---|---|--|
|                       | Size:<br>1000 µg/ml  | GenelD (NCBI):<br>3162                                  | Recommended Dilutions:<br>IF-P 1:50-1:500             |  |
|                       | Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG1190  | UNIPROT ID:<br>P09601                                   | Excitation/Emission maxima wavelengths:               |  |
|                       |  | Full Name:<br>heme oxygenase (decycling) 1              | 493 nm / 522 nm                                       |  |
|                       |  | Calculated MW:<br>33 kDa                                |   |  |
|                       |  | Observed MW:<br>28-33 kDa                               |   |  |
| Applications          | Tested Applications:   | Positive Controls:<br>IF-P : human liver cancer tissue, |   |  |
|                       | IF-P<br>Species Specificity:<br>human, mouse, rat  |   |   |  |
| Background Informatio | Heme oxygenase (HMOX1) catalyzes the first and rate-limiting step in the degradation of heme to yield equimolar quantities of biliverdin Ixa, carbon monoxide (CO), and iron. It has 3 isoforms: HO-1 is highly inducible, whereas HO-2 and HO-3 are constitutively expressed (PMID:10194478). Heme oxygenase-1 (HO-1) is expressed in many tissues and vascular smooth muscle cells, and endothelial cells (PMID:15451051) and has been identified as an important endogenous protective factor induced in many cell types by various stimulants, such as hemolysis, infiammatory cytokines,oxidative stress, heat shock, heavy metals, and endotoxin (PMID: 11522663). And the fullength HO-1 is very unstable and susceptible to truncation that generates an inactive, soluble form (28 kDa) (James R. Reed, Pharmacology, 535-568). |   |   |  |
| Storage               | Storage:<br>Store at -20°C. Avoid exposure to<br>Storage Buffer:<br>PBS with 50% Glycerol, 0.05% Pro<br>Aliquoting is unnecessary for -20  | oclin300, 0.5% BSA, pH 7.3.                             |   |  |

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite® Plus 488 HO-1/HMOX1 antibody (CL488-10701) at dilution of 1:200. Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite® Plus 488 HO-1/HMOX1 antibody (CL488-10701) at dilution of 1:200.