For Research Use Only

CoraLite® Plus 488-conjugated Ferritin light chain Polyclonal antibody




Catalog Number:CL488-10727 Featured Product			
Basic Information	Catalog Number: CL488-10727	GenBank Accession Number: BC008439	Purification Method: Antigen affinity purification
	Size: 1000 µg/ml	GenelD (NCBI): 2512	Recommended Dilutions: IF/ICC 1:50-1:500
	Source: Rabbit	UNIPROT ID: P02792	Excitation/Emission maxima wavelengths:
	lsotype: IgG	Full Name:493 nm / 522 nmferritin, light polypeptide	
	Immunogen Catalog Number: AG1149	Calculated MW: 19 kDa	
Applications	Tested Applications: IF/ICC Species Specificity: human, mouse, rat	Positive Controls: IF/ICC : HepG2 cells,	
Background Information	FTL belongs to the ferritin family. It stores iron in a soluble, non-toxic, readily available form. FTL is important for iron homeostasis. It plays a role in delivery of iron to cells. FTL mediates iron uptake in capsule cells of the developing kidney. Mutation of FTL will cause hereditary hyperferritinemia-cataract syndrome (HHCS) or neurodegeneration with brain iron accumulation type 3 (NBIA3). Ferritin light polypeptide (FTL) and ferritin heavy polypeptide (FTH1) were the main constituents the striatum and cerebellar cortex revealed.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 Ferritin light chain antibody (CL488-10727) at dilution of 1:200.